

SUPREP Bowel Prep Patient Instructions Split Dose Regimen

	cedure Date:
	cedure Location:val Time:
DAY	BEFORE PROCEDURE:
	NOT FOLLOW THE PREP DIRECTIONS ON THE BOX/ CONTAINER. FOLLOW THESE ONLY: escription has been sent electronically to
	You may have breakfast and lunch until 1:00pm then remain on clear liquids the rest of the day. Any of the following clear liquids are OK: Water
•	en amount pare (mane at pare) more and grape; or mine or an activity
•	collection to the collection of the collection o
9 TH	Jell-O (NO RED OR ORANGE) and Popsicles (NO RED OR ORANGE) ne evening before your colonoscopy beginning at <u>5:00 p.m.,</u> pour ONE (1) 6-ounce bottle of SUPREP
lic	juid into the mixing cup.
4. D	dd cool drinking water or sugar-free drink to the 16-ounce line on the cup and mix. rink ALL t he liquid in the mixing cup. There is no need to chug the mixture down, just drink it within the
5. Yo	our. Our MUST drink two (2) more 16-ounce cups fluid over the next hour or two. You may drink more fluids aroughout the night if you wish.
МОІ	RNING OF YOUR PROCEDURE:
	ne morning of your colonoscopy beginning at a.m. pour ONE (1) 6-ounce bottle of UPREP liquid into the mixing cup.
	dd cool drinking water or sugar-free drink to the 16-ounce line on the cup and mix. You must drink all e liquid in the next hour.
	ou MUST drink (2) more 16-ounce cups of fluids over the next 1 hour.
pr	ease note that you may not have anything to drink 4 hours prior to your procedure or your occedure may have to be rescheduled. You must be done with the entire prep prior to
qı	ead attached pages as to what medications to take prior to procedure. Please call with any uestions.
	ease have your driver with you at all times. You may not drive yourself home. ASE NOTE THAT YOU WILL BE CHARGED A CANCELLATION FEE IF YOU DO NOT CANCEL
PLE/	ASE NOTE THAT TOO WILL BE CHARGED A CANCELLATION FEETF TOO DO NOT CANCEL

YOUR PROCEDURE AS POLICY STATES. (Practice policy for administrative services.)



- 1. If you think that bowel prep did not completely clear the intestines for any reason, please call your doctor or the doctor on call for an alternative prep.
- 2. If you have a history of artificial heart valve, recent joint replacements or have been hospitalized for bacterial endocarditis, **please call your physician's nurse** so that antibiotic coverage can be arranged prior to your procedure.
- 3. If you are on any type of prescribed medications, **please notify the prescribing doctor** of your upcoming colonoscopy/surgery. Please make sure that doctor agrees that it is okay to stop the medication.
- 4. Do not take diuretics on the day of your procedure. These include clorothiazide, hydrochorothiazide, furosemide, bumetanide, toresmide, amiloride, spironoactone, triameterene, mannitol, dyazide, maxzide and moduretic.
- 5. Please make sure that you take all medications for acid reflux, asthma, bronchitis or emphysema in the morning with a sip of water on the day of your procedure.
- 6. If you take blood pressure medications in the morning, please make sure you take all blood pressure medications the day of your procedure with a sip of water. This includes combination medications for blood pressure which contain diuretics.

These are some examples of medications that should be stopped 3-7 days prior to your procedure. These medications thin the blood. If you are on a medication that was prescribed to you (or is over the counter) to help thin your blood and it is not on the list, please stop it prior to your procedure. (if recommended by your prescribing physician).

Blood thinners	NSAIDS	OTC Supplements
Aggrenox (dipyridamole+ aspirin)	Aspirin (Excedrin, bayer)	Fish oil
Aggrastat (tirofiban)	Celecoxib (celebrex)	Garlic in large amounts
Arixtra (fondaparinux sodium)	Ibuprofen (advil, motrin)	Ginko biloba
Coumadin (warfarin)	Indomethacin (indocin)	Ginger
Dicumarol (dicumarol)	Mag salicylate (bayer select)	Ginseng
Effient (prasugrel)	Meloxicam (mobic)	St John's Wart
Integrilin (eptifibatide)	Nabumetone (Relafen)	Saw Palmento
Miradon (anisinidine)	Naproxen (naprosyn)	Vitamin E
Plavix (clopidogrel bisulfate)	Naproxen sodium (aleve)	
Persantine (dipyridamole)	Oxaprozin (daypro)	
Pletal (cilostazol)	Piroxicam (feldene)	
Pradaxa (dabigatran)	Tolmetin sodium (tolectin)	
Reopro (abciximab)	Valdecoxib (bextra)	
Xeralto (rivaroxaban)		



ACE INIHIBITORS	BETA BLOCKERS	ANGIOTENSION II	COMBINATION
Accupril (quinapril)	Betapace (sotalol)	Atacand	Aldoril (methyldopa +
		(candesartan	hctz)
		cilexetil)	
Aceon (perindopril)	Bisoprolol (zebeta)	Avapro (irbesartan)	Capozide (captopril +
			hctz)
Altace (Ramipril)	Blocadren (timolol)	Benicar (olmesarten)	Corzide (nadolol +
			bendroflumethiazide)
Capoten (captopril)	Bystolic (nebivolol)	Cozaar (losartan)	Dyazide (triameterene +
			hctz)
Lotension (benazepril)	Cartrol (carteolol)	Diovan (Valsartan)	Hyzaar (losartan + hctz)
Mavik (trandolapril)	Coreg (carvedilol)	Hyzaar (losartan)	Inderide (propanolol +
			hctz)
Monopril (fosinopril)	Corgard (nadolol)	Micardis (telmisartan)	Lexxel(feldopine +
			enalapril)
Prinivil (lisinopril)	Inderal& inderal-la	Teveten (eprosartan	Lotrel (amlodipine +
	(propranolol)	mesylate)	benazepril)
Prinizide (lisinopril +	Kerlone(betaxolol)	Valsartan (diovan)	Lotension hct
diuretic)			(benazepril + hctz)
Univasc (moexipril)	Levatol (penbutolol)	OTHER BP MEDS	Moduretic (amiloride + hctz)
Vasotec (enalapril,	Lopressor (metoprolol)	Catapres (clonidine)	Prinizide (lisinopril +
analaprilat)	Lopiessoi (metopioloi)	Catapres (cionidine)	hctz)
Zesteril (lisinopril)	Normodyne(labetalol)	Cardura (doxazosin)	Tentoretic (atenolol +
, ,	, , ,	,	chlorthialidone)
	Tenormin (atenolol)	Cardene	Vaseretic(enalapril +
	,	(nicardipine)	diuretic)
	Toprol xl(metoprolol)	Cardizem (diltiazem)	Zestoretic (lisinopril +
		,	diurectic)
	Trandate (labetalol)	Norvasc (amlodipine)	Ziac (bisoprolol + hctz)
	Zebeta(bisoprolol)	Plendil (felodipine)	
		Procadia xl	
		(nifedipine)	