

Prepopik Bowel Prep for Afternoon Procedures

Procedure Date: _____

Procedure Location: _____

Arrival Time: _____

DAY BEFORE THE PROCEDURE: _____

DO NOT FOLLOW THE PREP DIRECTIONS ON THE BOX. FOLLOW THESE ONLY:

1. You may have a regular breakfast until 11:00am then remain on clear liquids the remainder of the day.

Any of the following clear liquids are OK:

- Water
- Strained fruit juices (without pulp) including apple, white grape, or white cranberry
- Gatorade
- Coffee or tea (DO NOT use any dairy or non-dairy creamer)
- Chicken broth and/ or Beef Broth
- Jell-o (NO RED OR PURPLE) and Popsicles (NO RED OR PURPLE)

First Dose:

A prescription has been sent electronically to your pharmacy _____.

STEP 1 - At 5:00 pm; you must fill the dosing cup provided with cold water to the lower (5 oz) line on the cup.

STEP 2 - Pour the contents of ONE packet and stir for 2-3 minutes until it is completely dissolved and drink the entire contents.

STEP 3- You must drink another five (5) 8-ounce cups of clear liquid, taken at your own pace within the next five (5) hours.

MORNING OF PROCEDURE: _____

Second Dose:

STEP 1- At _____, you must fill the dosing cup provided with cold water to the lower (5oz) line on the cup.

STEP 2 – Pour the contents of ONE (1) packet and stir for 2 minutes until it is completely dissolved and drink the entire contents.

STEP 3 – You must drink another three (3) 8- ounce cups of clear liquid before _____. Please note that you must *not have anything by mouth 4 hours prior to your procedure.*

- Nothing to eat or drink until after the procedure. This includes chewing gum, tobacco and candy.
- **Read attached pages as to what medications to take prior to procedure. Please call the office with any questions.**
- Bring insurance cards. Leave valuables at home.
- IT IS IMPERATIVE TO HAVE YOUR DRIVER HERE AT ALL TIMES. NO TAXIS ALLOWED!

PLEASE NOTE THAT YOU WILL BE CHARGED A CANCELLATION FEE IF YOU DO NOT CANCEL YOUR PROCEDURE AS POLICY STATES. (Practice policy for administrative service.)

1. If you think that your bowel prep did not completely clear the intestines for any reason, please call your doctor or the doctor on call for an alternative prep.
2. If you have a history of artificial heart valve, recent joint replacements or have been hospitalized for bacterial endocarditis, **please call your physician's nurse** so that antibiotic coverage can be arranged prior to your procedure.
3. If you are on any type of prescribed medications, **please notify the prescribing doctor** of your upcoming colonoscopy/surgery. Please make sure that doctor agrees that it is okay to stop the medication.
4. Do not take diuretics on the day of your procedure. These include clorothiazide, hydrochlorothiazide, furosemide, bumetanide, toresmide, amiloride, spironoactone, triameterene, mannitol, dyazide, maxzide and moduretic.
5. **Please make sure that you take all medications for acid reflux, asthma, bronchitis or emphysema in the morning with a sip of water on the day of your procedure.**
6. **If you take blood pressure medications in the morning, please make sure you take all blood pressure medications the day of your procedure with a sip of water. This includes combination medications for blood pressure which contain diuretics.**

These are some examples of medications that should be stopped 3-7 days prior to your procedure. These medications thin the blood. If you are on a medication that was prescribed to you (or is over the counter) to help thin your blood and it is not on the list, please stop it prior to your procedure. **(if recommended by your prescribing physician).**

Blood thinners	NSAIDS	OTC Supplements
Aggrenox (dipyridamole+ aspirin)	Aspirin (Excedrin, bayer)	Fish oil
Aggrastat (tirofiban)	Celecoxib (celebrex)	Garlic in large amounts
Arixtra (fondaparinux sodium)	Ibuprofen (advil, motrin)	Ginko biloba
Coumadin (warfarin)	Indomethacin (indocin)	Ginger
Dicumarol (dicumarol)	Mag salicylate (bayer select)	Ginseng
Effient (prasugrel)	Meloxicam (mobic)	St John's Wart
Integrilin (eptifibatide)	Nabumetone (Relafen)	Saw Palmento
Miradon (anisinidine)	Naproxen (naprosyn)	Vitamin E
Plavix (clopidogrel bisulfate)	Naproxen sodium (aleve)	
Persantine (dipyridamole)	Oxaprozin (daypro)	
Pletal (cilostazol)	Piroxicam (feldene)	
Pradaxa (dabigatran)	Tolmetin sodium (tolectin)	
Reopro (abciximab)	Valdecoxib (bextra)	
Xeralto (rivaroxaban)		

ACE INIHIATORS	BETA BLOCKERS	ANGIOTENSION II	COMBINATION
Accupril (quinapril)	Betapace (sotalol)	Atacand (candesartan cilexetil)	Aldoril (methyldopa + hctz)
Aceon (perindopril)	Bisoprolol (zebeta)	Avapro (irbesartan)	Capozide (captopril + hctz)
Altace (Ramipril)	Blocadren (timolol)	Benicar (olmesarten)	Corzide (nadolol + bendroflumethiazide)
Capoten (captopril)	Bystolic (nebivolol)	Cozaar (losartan)	Dyazide (triameterene + hctz)
Lotension (benazepril)	Cartrol (carteolol)	Diovan (Valsartan)	Hyzaar (losartan + hctz)
Mavik (trandolapril)	Coreg (carvedilol)	Hyzaar (losartan)	Inderide (propranolol + hctz)
Monopril (fosinopril)	Corgard (nadolol)	Micardis (telmisartan)	Lexxel(feldopine + enalapril)
Prinivil (lisinopril)	Inderal& inderal-la (propranolol)	Teveten (eprosartan mesylate)	Lotrel (amlodipine + benazepril)
Prinizide (lisinopril + diuretic)	Kerlone(betaxolol)	Valsartan (diovan)	Lotension hct (benazepril + hctz)
Univasc (moexipril)	Levatol (penbutolol)	OTHER BP MEDS	Moduretic (amiloride + hctz)
Vasotec (enalapril, analaprilat)	Lopressor (metoprolol)	Catapres (clonidine)	Prinizide (lisinopril + hctz)
Zesteril (lisinopril)	Normodyne(labetalol)	Cardura (doxazosin)	Tentoretic (atenolol + chlorthialidone)
	Tenormin (atenolol)	Cardene (nicardipine)	Vaseretic(enalapril + diuretic)
	Toprol xl(metoprolol)	Cardizem (diltiazem)	Zestoretic (lisinopril + diurectic)
	Trandate (labetalol)	Norvasc (amlodipine)	Ziac (bisoprolol + hctz)
	Zebeta(bisoprolol)	Plendil (felodipine)	
		Procardia xl (nifedipine)	