



### Prepopik Bowel Prep for Morning Procedures

Facility: \_\_\_\_\_  
 Procedure Date:/Time: \_\_\_\_\_  
 Report to:/Report Time: \_\_\_\_\_  
 DAY BEFORE PROCEDURE: \_\_\_\_\_

#### PREPARATION DIET

The day before your colonoscopy you may have breakfast until 11am

After 11:00am you will be on a clear liquid diet such as: **water**, **clear fruit juices** (ex: apple juice, white grape juice, white cranberry juice), **Gatorade®** (except purple, orange and red flavors/tints), **clear broth** (chicken or beef), **Jell-o®** (except purple, orange and red flavors/tints), **weak tea** (without cream), **coffee** (without cream), **icy Popsicles** (except purple, orange and red flavors/tints), and **regular or diet soft drinks** (except orange, purple and red flavors/tints). *(A clear liquid is anything that you can read the newspaper through. THIS DOES NOT INCLUDE ALCOHOL)*

**PLEASE FOLLOW THE PREPERATION INSTRUCTIONS GIVEN TO YOU BY OUR OFFICE, NOT THE ONES ON THE CONTAINER/BOX**

**At 3:00pm** – Prepare the first 5 ounces of solution using a packet of Prepopik:  
 Tear open one of the packets provided in the kit and pour it into the container. Fill the container with water to the 5 ounce mark. Stir until powder has dissolved. **After drinking the five ounce of Prepopik solution**, drink no less than an additional **five 8-ounce** (the upper line on the dosing cup) cups of clear liquids at a rate that is comfortable for you.

**At 9:00pm** – Prepare the second 5 ounces of solution the same as you did the first. **After drinking the five ounce of Prepopik solution**, drink no less than an additional **three 8-ounce** of clear liquids at a rate that is comfortable for you. You may continue to drink clear liquids until midnight.

**NOTHING TO EAT OR DRINK AFTER MIDNIGHT THE DAY BEFORE YOUR PROCEDURE. THIS INCLUDES WATER, CHEWING GUM, HARD OR SOFT CANDY ETC.**

**MORNING of PROCEDURE:** \_\_\_\_\_

1. Bring insurance card(s). Leave valuables at home.
2. Please note that all blood pressure medications should be taken the morning/evening prior to your surgery with a sip of water. Do not take any diuretics prior to surgery.

**SURGICAL OR ENDOSCOPY PROCEDURES CANCELLED AND RESCHEDULED WITH LESS THAN 3 BUSINESS DAYS NOTICE WILL INCUR A \$100 FEE.**

**NO DRIVING UP TO 12 HOURS AFTER YOUR PROCEDURE**  
*Due to the sedation given for this procedure, you will need a designated driver. It is imperative that your driver remains in the waiting area at all times.*

1. If you think that your bowel prep did not completely clear the intestines for any reason, please call your doctor or the doctor on call for an alternative prep.
2. If you have a history of artificial heart valve, recent joint replacements or have been hospitalized for bacterial endocarditis, **please call your physician's nurse** so that antibiotic coverage can be arranged prior to your procedure.
3. If you are on any type of prescribed medications, **please notify the prescribing doctor** of your upcoming colonoscopy/surgery. Please make sure that doctor agrees that it is okay to stop the medication.
4. Do not take diuretics on the day of your procedure. These include clorothiazide, hydrochlorothiazide, furosemide, bumetanide, toresmide, amiloride, spironoactone, triameterene, mannitol, dyazide, maxzide and moduretic.
5. **Please make sure that you take all medications for acid reflux, asthma, bronchitis or emphysema in the morning with a sip of water on the day of your procedure.**
6. **If you take blood pressure medications in the morning, please make sure you take all blood pressure medications the day of your procedure with a sip of water. This includes combination medications for blood pressure which contain diuretics.**

These are some examples of medications that should be stopped 3-7 days prior to your procedure. These medications thin the blood. If you are on a medication that was prescribed to you (or is over the counter) to help thin your blood and it is not on the list, please stop it prior to your procedure. (if recommended by your prescribing physician).

Blood thinners	NSAIDS	OTC Supplements
<b>Aggrenox (dipyridamole+ aspirin)</b>	<b>Aspirin ( Excedrin, bayer)</b>	<b>Fish oil</b>
<b>Aggrastat (tirofiban)</b>	<b>Celecoxib (celebrex)</b>	<b>Garlic in large amounts</b>
<b>Arixtra (fondaparinux sodium)</b>	<b>Ibuprofen (advil, motrin)</b>	<b>Ginko biloba</b>
<b>Coumadin (warfarin)</b>	<b>Indomethacin ( indocin)</b>	<b>Ginger</b>
<b>Dicumarol (dicumarol)</b>	<b>Mag salicylate (bayer select)</b>	<b>Ginseng</b>
<b>Effient (prasugrel)</b>	<b>Meloxicam (mobic)</b>	<b>St John's Wart</b>
<b>Integrilin (eptifibatide)</b>	<b>Nabumetone ( Relafen)</b>	<b>Saw Palmento</b>
<b>Miradon ( anisinidine)</b>	<b>Naproxen (naprosyn)</b>	<b>Vitamin E</b>
<b>Plavix (clopidogrel bisulfate)</b>	<b>Naproxen sodium (aleve)</b>	
<b>Persantine (dipyridamole)</b>	<b>Oxaprozin (daypro)</b>	



<b>Pletal (cilostazol)</b>	<b>Piroxicam (feldene)</b>	
<b>Pradaxa (dabigatran)</b>	<b>Tolmetin sodium (tolectin)</b>	
<b>Reopro (abciximab)</b>	<b>Valdecoxib (bextra)</b>	
<b>Xeralto (rivaroxaban)</b>		

<b>ACE INHIBITORS</b>	<b>BETA BLOCKERS</b>	<b>ANGIOTENSION II</b>	<b>COMBINATION</b>
Accupril (quinapril)	Betapace (sotalol)	Atacand (candesartan cilexetil)	Aldoril (methyldopa + hctz)
Aceon ( perindopril)	Bisoprolol (zebeta)	Avapro (irbesartan)	Capozide (captopril + hctz)
Altace (Ramipril)	Blocadren (timolol)	Benicar (olmesartan)	Corzide (nadolol + bendroflumethiazide)
Capoten (captopril)	Bystolic (nebivolol)	Cozaar (losartan)	Dyazide (triameterene + hctz)
Lotension (benazepril)	Cartrol (carteolol)	Diovan (Valsartan)	Hyzaar (losartan + hctz)
Mavik (trandolapril)	Coreg (carvedilol)	Hyzaar (losartan)	Inderide (propranolol + hctz)
Monopril (fosinopril)	Corgard (nadolol)	Micardis (telmisartan)	Lexxel( feldopine + enalapril)
Prinivil (lisinopril)	Inderal& nderal-la (propranolol)	Teveten (eprosartan mesylate)	Lotrel (amlodipine + benazepril)
Prinizide (lisinopril + diuretic)	Kerlone(betaxolol)	Valsartan (diovan)	Lotension hct (benazepril + hctz)
Univasc (moexipril)	Levatol (penbutolol)	<b>OTHER BP MEDS</b>	Moduretic (amiloride + hctz)
Vasotec (enalapril, analaprilat)	Lopressor (metoprolol)	Catapres (clonidine)	Prinizide (lisinopril + hctz)
Zesteril (lisinopril)	Normodyne(labetalol)	Cardura (doxazosin)	Tentoretic (atenolol + chlorthalidone)
	Tenormin (atenolol)	Cardene (nicardipine)	Vaseretic(enalapril + diuretic)
	Toprol xl(metoprolol)	Cardizem (diltiazem)	Zestoretic (lisinopril + diurectic)
	Trandate (labetalol)	Norvasc (amlodipine)	Ziac (bisoprolol + hctz)
	Zebeta(bisoprolol)	Plendil (felodipine)	
		Procardia xl (nifedipine)	